Gynecomastia Preoperative Instructions

Υ (our day/time of surgery is The location is
•	Please arrive to the facility at least one hour prior to your scheduled time.
•	If you are a smoker, please stop for a minimum of 4 weeks prior to surgery. Smoking reduces circulation to the skin and will slow your healing.
•	Avoid Aspirin, Ibuprofen, Advil, Motrin, Naprosyn, Aleve, Celebrex, Excedrin, Fish Oil and Vitamin E containing products for 2 weeks prior to surgery. This also includes the use of multivitamins and herbal supplements. All of these medications have a tendency to prolong bleeding. Please review the attached sheet to confirm the use of any of these. <i>Tylenol is preferred for pain relief.</i>
•	Please purchase at the pharmacy a tube of antibiotic ointment (such as Bacitracin or Neosporin), gauze sponges, and surgical tape (preferably paper tape).
•	You have received prescriptions for You should have these filled <i>prior</i> to your day of surgery. You will not need to take any of them until after surgery.
•	You should have your bloodwork drawn at least two weeks prior to your surgery.
•	Shower with an antibacterial soap the day before or the morning of your surgery.
•	Have NOTHING to eat or drink after midnight on the night before your surgery.
•	You will be provided with a compressive garment, which you will wear for one month following your surgery.
•	We recommend the use of silicone sheeting for scar therapy. This product can be purchased at CVS pharmacy. You will start using this or any type of scar therapy approximately 2 weeks after surgery.
•	On the day of surgery, wear a large, front closure shirt and loose fitting pants with an elastic waist.
Ot	her: