Labiaplasty Preoperative Instructions

Y	our day/time of surgery is	. The location is	
•	Please arrive to the facility at least one hour prior to your scheduled time.		
•	If you are a smoker, please stop for a minimum of 4 weeks prior to surgery. Smoking reduces circulation to the skin and will slow your healing.		
•	Avoid Aspirin, Ibuprofen, Advil, Motrin, Naprosyn, Aleve, Celebrex, Excedrin, Fish Oil and Vitamin E containing products for 2 weeks prior to surgery. This also includes the use of multivitamins and herbal supplements. All of these medications have a tendency to prolong bleeding. Please review the attached sheet to confirm the use of any of these. <i>Tylenol is preferred for pain relief</i> .		
•	You have received prescriptions for You should have these filled <i>prior</i> to your day of surgery. You will not need to take any of them until after surgery.		
•	You should have your bloodwork drawn at least two weeks prior to your surgery.		
•	Have NOTHING to eat or drink after midnight on the night before your surgery.		
•	Please purchase antibiotic ointment (such as Bacitracin or Neosporin) for postoperative wound care and a perineal irrigation bottle for post void cleansing. We also recommend using Dermoplast Antibacterial Pain Relieving Spray and TUCKS Medicated pads for postoperative discomfort. These items can be purchased over-the-counter at your local pharmacy or online at www.amazon.com.		
•	On the day of surgery, wear a large, front of	closure shirt and loose fitting pants with an elastic waist.	
O1	Other:		