## **Medications & Supplements to Avoid**

The following medications, vitamins, and herbal supplements can interfere with your surgery and/or recovery. Some may increase bleeding, bruising, or risk of blood clots. Please check all or that you take occasionally or on a regular basis. \*Note this also includes the use of any diet/weight loss pills\*

## Please stop all two weeks prior to surgery until one week after surgery:

**Estrogen medications:** 

**ALL nicotine products!** 

Anorex

CaroExpel

Centramine

Cortislim

**Ephedra** 

Leptopril

Relacore

HoodiElite

Phentermine

Birth control pills Hormone replacement

**Anticoagulants:** 

Coumadin Arixtra Lovenox

Plavix Xarelto

**Aspirin:** ASA all forms

Bufferin Excedrin

**Anti-inflammatories:** 

Aleve Ibuprofen Midol Celebrex Advil Naproxen Motrin **Pamprin** Excedrin Naprosyn Diclofenac Meloxicam

**Vitamins containing >** 400 units of vitamin E

**Vitamin** C (more than 2000mg daily)

Any herbal medications: **Diet Pills:** 

> Aloe Vera Gingko Biloba Bromelain Ginseng Goldenseal Danshen Dong Quai Kava Echinacea Licorice **Ephedra** Omega-3 Feverfew Senna

Flax Seed St. John's Wort (all types)

Fish Oils Valerian

Slim Citi Hoodia Garlic Xenadrine Ginger

All cold medications and decongestants